

Goncharov's programme, and depth reductions of multiple polylogarithms

Steven Charlton
Max Planck Institute for Mathematics, Bonn

21 June 2024
 Spec($\overline{\mathbb{Q}}(2\pi i)$), Fields Institute, Toronto

Multiple polylogarithm definitions

Definition

Multiple polylogarithm (MPL) is

$$\text{Li}_{k_1, k_2, \dots, k_d}(x_1, x_2, \dots, x_d) := \sum_{0 < m_1 < m_2 < \dots < m_d} \frac{x_1^{m_1} \cdots x_d^{m_d}}{m_1^{k_1} \cdots m_d^{k_d}}, \quad |x_i| < 1.$$

Generalises

$$\text{Li}_1(x) = -\log(1-x).$$

- **Weight** is $n = k_1 + \cdots + k_d$,
- **Depth** is d .

Goal: Fundamental challenge around MPL's is to understand the filtration by depth.

Relating different depths

Functions of different depths can be related:

$$\text{Li}_1(x) \text{Li}_1(y) = \text{Li}_{1,1}(x, y) + \text{Li}_{1,1}(y, x) + \text{Li}_2(xy)$$

$$\leftrightarrow \sum_{m=1}^{\infty} \sum_{\ell=1}^{\infty} \frac{x^m y^\ell}{m \cdot \ell} = \left(\sum_{m<\ell} + \sum_{m>\ell} + \sum_{m=\ell} \right) \frac{x^m y^\ell}{m \cdot \ell}$$

So-called **shuffle product** of MPL's \rightsquigarrow this generalises easily

More surprising when the depth reduces

Proposition (Goncharov, Zagier, Lewin^(?))

As a power-series identity, for $|xy| < 1, |y| < 1$,

$$\text{Li}_{1,1}(x, y) = \text{Li}_2\left(\frac{y(x-1)}{1-y}\right) - \text{Li}_2\left(\frac{-y}{1-y}\right) - \text{Li}_2(xy).$$

Further depth reductions?

Proposition ($\text{Li}_{1,1,1}$ is depth 1, Goncharov, Zhao, Lewin^(?), ...)

$$\begin{aligned} \text{Li}_{1,1,1}(x, y, z) = & -\text{Li}_3\left(\frac{1-xyz}{1-x}\right) - \text{Li}_3\left(\frac{1-xyz}{xy(1-z)}\right) + \text{Li}_3\left(\frac{(y-1)(1-xyz)}{(1-x)y(1-z)}\right) + \text{Li}_3(xy) \\ & - \text{Li}_3\left(\frac{y(1-x)}{y-1}\right) + \text{Li}_3(1-x) - \text{Li}_3\left(\frac{y(1-z)}{y-1}\right) + \text{Li}_3\left(\frac{-y}{1-y}\right) + \text{products} \end{aligned}$$

Can weight 4 be reduced to depth 1? Apparently not, but:

Proposition ($\text{Li}_{3,1}$ relations, mod depth 1, Zagier, Gangl)

$$\begin{aligned} \text{Li}_{3,1}\left(\frac{1-x}{y}, y\right) + \text{Li}_{3,1}\left(\frac{x}{y}, y\right) = & -\frac{1}{2} \text{Li}_4\left(\frac{(1-x)y}{x(1-y)}\right) - \frac{1}{2} \text{Li}_4\left(\frac{xy}{(1-x)(1-y)}\right) + \frac{1}{2} \text{Li}_4\left(\frac{(1-y)y}{(1-x)x}\right) \\ & - \text{Li}_4\left(\frac{1-y}{1-x}\right) - \text{Li}_4\left(\frac{1-y}{x}\right) + \text{Li}_4\left(-\frac{y}{1-y}\right) \\ & - \text{Li}_4(1-x) - \text{Li}_4(x) + 2 \text{Li}_4(y) + \text{products} \end{aligned}$$

Question: How to predict or understand, such reductions and obstructions?

Connections elsewhere

Values of zeta functions: [[see Kairi Black's lightning talk, Tuesday]]

- Dedekind zeta function of a number field F is $\zeta_F(s) = \sum_{I \neq 0 \subset \mathcal{O}_F} N(I)^{-s}$.
- Analytic class number formula: $\text{Res}_{s=1} \zeta_F(s)$ via $\underbrace{\log / \text{Li}_1}_{\text{of units}} \text{ regulator } R_F$ (& arithmetic data).
- Zagier: conjecture for $\zeta_F(n)$ via Li_n (& 'higher units' in K -theory).
- Goncharov relates $\zeta_F(n)$ to high-depth "Grassmannian polylogarithm".

Task: How to reduce to depth 1?

Also: hyperbolic geometry, cluster algebras, high-energy physics, ...

Motivic multiple polylogarithms

Recall: Iterated integral [[special case from Rosa Preiß's talk, Thursday]]

$$I_{(\gamma)}(a; x_1, \dots, x_n; b) := \int_{a < t_1 < \dots < t_n < b} \frac{dt_1}{t_1 - x_1} \wedge \frac{dt_2}{t_2 - x_2} \wedge \dots \wedge \frac{dt_n}{t_n - x_n}.$$

By term-wise integration of geometric series (with $X_j := \prod_{i=1}^j x_i$),

$$\text{Li}_{k_1, k_2, \dots, k_d}(x_1, \dots, x_d) = (-1)^d I(0; 1, \underbrace{0, \dots, 0}_{k_1}, X_1, \underbrace{0, \dots, 0}_{k_2}, X_2, \dots, X_{d-1}, \underbrace{0, \dots, 0}_{k_d}; X_d).$$

- Goncharov: upgrade of I to framed mixed Tate motives $I^{\mathfrak{u}}(a; x_1, \dots, x_n; b)$,
 $I^{\mathfrak{u}}$ lives in connected, weight graded Hopf algebra $(\mathcal{A}_\bullet(\overline{\mathbb{Q}}), \Delta)$


Idea: Can keep track of differential form and integration chain separately

- Define $\text{Li}_{k_1, k_2, \dots, k_d}^{\mathfrak{u}}(x_1, \dots, x_d)$ via $I^{\mathfrak{u}}$

Lie coalgebra, cobracket, depth filtration

- $\mathcal{L}_\bullet = \mathcal{A}_\bullet / \mathcal{A}_{>0}^2$ = motivic MPL's modulo products, obtain $\Delta: \mathcal{L}_\bullet(F) \rightarrow \bigwedge^2 \mathcal{L}_\bullet(F)$

$$\Delta I^\mathcal{L}(x_0; x_1, \dots, x_n; x_{n+1}) = \sum_{i < j} I^\mathcal{L}(x_i; x_{i+1}, \dots, x_{j-1}; x_j) \wedge I^\mathcal{L}(x_0; x_1, \dots, x_i, x_j, \dots, x_n; x_{n+1})$$



- Depth filtration $\mathcal{D}_d = \{\text{depth } \leq d \text{ MPL's in } \mathcal{L}_\bullet\}$ has

$$\Delta(\mathcal{D}_d) \subset \mathcal{D}_d \wedge \mathcal{L}_1 \oplus \bigoplus_{\substack{i+j=d \\ i,j \geq 1}} \mathcal{D}_i \wedge \mathcal{D}_j.$$

Why? Follows since $I^\mathcal{L}(a; \overbrace{0, \dots, 0}^\ell; b) = \frac{1}{\ell!} I^\mathcal{L}(a; 0, b)^\ell \equiv 0 \pmod{\text{products}}$

Cobrackets and obstructions to depth 1

$$\Delta \text{Li}_4^{\mathcal{L}}(x) = \log^{\mathcal{L}}(x) \wedge \text{Li}_3^{\mathcal{L}}(x) \in \mathcal{L}_1 \wedge \mathcal{L}_3,$$

$$\begin{aligned} \Delta \text{Li}_{3,1}^{\mathcal{L}}(x, y) = & -\log^{\mathcal{L}}(x) \wedge \text{Li}_{2,1}^{\mathcal{L}}(x, y) + \text{Li}_1^{\mathcal{L}}(y) \wedge \text{Li}_3^{\mathcal{L}}(x) - \text{Li}_1^{\mathcal{L}}(y) \wedge \text{Li}_3^{\mathcal{L}}(xy) \\ & - \text{Li}_1^{\mathcal{L}}(xy) \wedge \text{Li}_3^{\mathcal{L}}(x) + \text{Li}_1^{\mathcal{L}}(xy) \wedge \text{Li}_3^{\mathcal{L}}(y) \\ & + \text{Li}_2^{\mathcal{L}}(y) \wedge \text{Li}_2^{\mathcal{L}}(xy) \end{aligned} \quad \left. \right\} \in \mathcal{L}_1 \wedge \mathcal{L}_3 \in \mathcal{L}_2 \wedge \mathcal{L}_2$$

- Write $\overline{\Delta}$ to mean ignore \mathcal{L}_1 , then we obtain

$$\overline{\Delta} \text{Li}_4^{\mathcal{L}}(x) = 0, \quad \overline{\Delta} \text{Li}_{3,1}^{\mathcal{L}}(x, y) = \text{Li}_2^{\mathcal{L}}(y) \wedge \text{Li}_2^{\mathcal{L}}(xy).$$

Hence $\text{Li}_{3,1}^{\mathcal{L}}(x, y) \neq \sum \text{Li}_4^{\mathcal{L}}$'s.

Question: Is this the only obstruction?

Goncharov Depth Conjecture (depth 1, simplified version)

A linear combination of MPL's has depth 1 if and only if $\overline{\Delta}$ vanishes.

Expectations and evidence

Weight 2, 3: No space for non-trivial contribution, so

$$\overline{\Delta}(\mathcal{L}_2) = 0, \quad \overline{\Delta}(\mathcal{L}_3) = 0.$$

Expect $\text{Li}_{1,1}^{\mathcal{L}}(x, y) = \sum \text{Li}_2^{\mathcal{L}} \cdot s$, and $\text{Li}_{1,1,1}^{\mathcal{L}}(x, y, z) = \sum \text{Li}_3^{\mathcal{L}} \cdot s$. ✓ (Intro)

Weight 4: After tweaking variables $\overline{\Delta} \text{Li}_{3,1}^{\mathcal{L}}\left(\frac{x}{y}, y\right) = -\text{Li}_2^{\mathcal{L}}(x) \wedge \text{Li}_3^{\mathcal{L}}(y)$.

Vanishing of $\overline{\Delta}\left(\sum_i \lambda_i \text{Li}_{3,1}^{\mathcal{L}}\left(\frac{x_i}{y}, y\right)\right)$ under dilogarithm identities $\sum_i \lambda_i \text{Li}_2(x_i) = 0$.

- Symmetries $\text{Li}_2^{\mathcal{L}}(x) + \text{Li}_2^{\mathcal{L}}(1-x) = 0$ and $\text{Li}_2^{\mathcal{L}}(x) + \text{Li}_2^{\mathcal{L}}(x^{-1}) = 0$ ✓ (Intro + ε)
- 5-term relation $\sum_{i=0}^4 (-1)^i \text{Li}_2^{\mathcal{L}}([z_0, \dots, \hat{z}_i, \dots, z_4]) = 0$, $[z_1, \dots, z_4] = \frac{(z_1-z_2)(z_3-z_4)}{(z_2-z_3)(z_4-z_1)}$

Five-term & cluster algebras [see José Simental's talk, Thursday]

• Example

$$\mathcal{F} = \mathbb{C}(x_1, x_2)$$

$$A \cong \mathbb{C}[x_1, x_2, x_3, x_5] / \left(\begin{array}{l} x_1 x_3 = 1 + x_2 \\ x_2 x_5 = 1 + x_1 \end{array} \right)$$

$$C = \{ \dots, x_{-2}, x_{-1}, x_0, \boxed{x_1}, \boxed{x_2}, \boxed{x_3}, \boxed{x_4}, x_5, \dots \} = \{x_1, x_2, x_3, x_4, x_5\}$$

Clusters: $\{x_i, x_{i+1}\}$, $i \in \mathbb{Z}$.

Mutation: $x_i x_{i+2} = 1 + x_{i+1}$

$$x_1, x_2, x_3 = \frac{1+x_2}{x_1}, \quad x_4 = \frac{1+x_3}{x_2} = \frac{1 + \frac{1+x_2}{x_1}}{x_2} = \frac{x_1 + x_2 + 1}{x_1 x_2}$$

$$x_5 = \frac{1+x_4}{x_3} = \frac{1 + \frac{x_1 + x_2 + 1}{x_1 x_2}}{\frac{1+x_2}{x_1}} = \frac{(x_1 x_2 + x_1 + x_2 + 1) x_1}{(1+x_2) x_1 x_2} = \frac{1+x_1}{x_2}$$

$$x_6 = \frac{1+x_5}{x_4} = \frac{1 + \frac{1+x_1}{x_2}}{\frac{x_1 + x_2 + 1}{x_1 x_2}} = \frac{(x_1 + x_2 + 1) x_1 x_2}{(x_1 + x_2 + 1) x_2} = x_1 \quad x_7 = x_2$$

Cluster algebra magic: $\sum_{i=1}^5 \text{Li}_2^{\mathcal{L}}(-x_i) = 0$

Expectations and evidence

Weight 2, 3: No space for non-trivial contribution, so

$$\overline{\Delta}(\mathcal{L}_2) = 0, \quad \overline{\Delta}(\mathcal{L}_3) = 0.$$

Expect $\text{Li}_{1,1}^{\mathcal{L}}(x, y) = \sum \text{Li}_2^{\mathcal{L}}\text{'s}$, and $\text{Li}_{1,1,1}^{\mathcal{L}}(x, y, z) = \sum \text{Li}_3^{\mathcal{L}}\text{'s}$. ✓ (Intro)

Weight 4: After tweaking variables $\overline{\Delta} \text{Li}_{3,1}^{\mathcal{L}}\left(\frac{x}{y}, y\right) = -\text{Li}_2^{\mathcal{L}}(x) \wedge \text{Li}_3^{\mathcal{L}}(y)$.

Vanishing of $\overline{\Delta}\left(\sum_i \lambda_i \text{Li}_{3,1}^{\mathcal{L}}\left(\frac{x_i}{y}, y\right)\right)$ under dilogarithm identities $\sum_i \lambda_i \text{Li}_2(x_i) = 0$.

- Symmetries $\text{Li}_2^{\mathcal{L}}(x) + \text{Li}_2^{\mathcal{L}}(1-x) = 0$ and $\text{Li}_2^{\mathcal{L}}(x) + \text{Li}_2^{\mathcal{L}}(x^{-1}) = 0$ ✓ (Intro + ε)
- 5-term relation $\sum_{i=0}^4 (-1)^i \text{Li}_2^{\mathcal{L}}([z_0, \dots, \widehat{z}_i, \dots, z_4]) = 0, \quad [z_1, \dots, z_4] = \frac{(z_1-z_2)(z_3-z_4)}{(z_2-z_3)(z_4-z_1)}$

Theorem (5-term reduction, Gangl 2012; Goncharov-Rudenko, Matveiakin-Rudenko)

$$\sum_{i=0}^4 (-1)^i \text{Li}_{3,1}^{\mathcal{L}}\left(\frac{\text{cr}(z_0, \dots, \widehat{z}_i, \dots, z_4)}{y}, y\right) = \sum_{i=1}^{122+\varepsilon} \text{Li}_4^{\mathcal{L}}(g_i(z_0, \dots, z_4, y)), \quad g_i \text{ rational}$$

Remark: Crucial ingredient to Goncharov-Rudenko proof of Zagier's Conjecture on $\zeta_F(4)$.

Iterating to understand higher depth

- Computing $\overline{\Delta}$ gets complicated:

$$\begin{aligned}\overline{\Delta} \text{Li}_{4,1,1}^{\mathcal{L}}(x, y, z) = & \text{Li}_2^{\mathcal{L}}(y) \wedge \text{Li}_{3,1}^{\mathcal{L}}(xy, z) - \text{Li}_{3,1}^{\mathcal{L}}\left(yz, \frac{1}{y}\right) \wedge \text{Li}_2^{\mathcal{L}}(xyz) - \text{Li}_3^{\mathcal{L}}(y) \wedge \text{Li}_{2,1}^{\mathcal{L}}(xy, z) \\ & - \text{Li}_{2,1}^{\mathcal{L}}\left(yz, \frac{1}{y}\right) \wedge \text{Li}_3^{\mathcal{L}}(xyz) + \text{Li}_4^{\mathcal{L}}\text{'s} \wedge \text{weight } 2 + \text{Li}_n^{\mathcal{L}}\text{'s} \wedge \text{Li}_m^{\mathcal{L}}\text{'s}.\end{aligned}$$

- Use $\overline{\Delta}$ to pick “genuine” depth 2 terms, so iterate¹ $\overline{\Delta}^{[2]} := \overline{\Delta} \circ (\overline{\Delta} \otimes \text{id})$

$$\overline{\Delta}^{[2]} \text{Li}_{4,1,1}^{\mathcal{L}}(x, y, z) = \text{Li}_2^{\mathcal{L}}(y) \otimes \text{Li}_2^{\mathcal{L}}(z) \otimes \text{Li}_2^{\mathcal{L}}(xyz) \quad (\text{in CoLie}_3).$$

Since $\overline{\Delta}^{[2]} \text{Li}_{a,b}^{\mathcal{L}}(x, y) = 0$, we deduce $\text{Li}_{4,1,1}^{\mathcal{L}}(x, y, z) \neq \sum \text{depth } 2$.

Question: Is this the only obstruction?

Goncharov Depth Conjecture (simplified version)

A linear combination of MPL's has depth $< k$ if and only if $\overline{\Delta}^{[k-1]}$ vanishes.

¹needs to be formalised: in a coLie algebra, the coJacobi identity would imply naïve $\Delta^2 = 0$

Expectations and evidence (II)

Observe: If $k > \frac{n}{2}$, then $\overline{\Delta}^{[k-1]}(\mathcal{L}_n) = 0$ (No space for k -many weight 2 pieces.)

Theorem (Rudenko, 2022)

Every weight n multiple polylogarithm can be expressed via depth $\leq \lfloor \frac{n}{2} \rfloor$ functions.

Idea: Explicit formula via “quadrangular” polylogs with depth $\frac{n}{2}$

Write:

$$f(x, y, z) := \text{Li}_{4,1,1}^{\mathcal{L}}((xyz)^{-1}, y, z)$$

$$\overline{\Delta}^{[2]} f(x, y, z) = -\text{Li}_2^{\mathcal{L}}(x) \otimes \text{Li}_2^{\mathcal{L}}(y) \otimes \text{Li}_2^{\mathcal{L}}(z) \quad (\text{in CoLie}_k)$$

Then $f(x, y, z)$ behaves like $\text{Li}_2^{\mathcal{L}}$ in each argument. Expect the following reductions:

- $f(x, y, z) + f(\underbrace{1-x, y, z}_{\text{or } x^{-1}}) \stackrel{?}{\equiv} 0 \pmod{\text{dp} \leq 2}$ (Zagier-type)
- $f(\text{dilog 5-term}, y, z) \stackrel{?}{\equiv} 0 \pmod{\text{dp} \leq 2}$ (Gangl-type)

Weight 6 results

Saw similar expectations hold in weight 4. They are also recently known in weight 6.

Write $f(x, y, z) := \text{Li}_{4,1,1}^{\mathcal{L}}((xyz)^{-1}, y, z)$, as before

Theorem (Gangl-type reduction, Matveiakin-Rudenko 2022)

$$f(\text{dilog 5-term}, y, z) \equiv 0 \pmod{\text{depth} \leq 2 \text{ \& the Zagier reductions}}$$

Theorem (Zagier-type reductions, C 2024, arXiv:2405.13853)

$$f(x, y, z) + f(1 - x, y, z) \equiv 0 \pmod{\text{depth} \leq 2}$$

$$f(x, y, z) + f(x^{-1}, y, z) \equiv 0 \pmod{\text{depth} \leq 2}$$

Consequence: Goncharov's Depth Conjecture holds for weight 6 depth 3 ($k = 3$).

(Zagier-reductions seem more difficult than the Gangl-reduction in general!)

Remark: Similar conjectural expectations in all even weights.

Ingredients for the proof – Quadrangular polylogarithms

Rudenko's quadrangular polylogarithms $\text{QLi}_{n+k}(x_0, \dots, x_{2n+1})$

- “Nice” MPL functions on $\mathfrak{M}_{0,2n+2}$, expressible via “quadrangulations” of polygons
- Weight-independent depth n expression via cross-ratio $[x_a, x_b, x_c, x_d] = \frac{(x_a-x_b)(x_c-x_d)}{(x_b-x_c)(x_d-x_a)}$

$$\begin{aligned} \text{QLi}_{k+2}(x_1, \dots, x_6) &= \text{Li}_{k;1,1}^{\mathcal{L}} \left(+ \begin{array}{c} x_2 \\ \diagdown \\ \text{hexagon} \\ \diagup \\ x_1 \\ - \end{array} \right. \\ &\quad \left. - \begin{array}{c} x_2 \\ \diagdown \\ \text{hexagon} \\ \diagup \\ x_1 \\ + \end{array} \right. \begin{array}{c} x_2 \\ \diagdown \\ \text{hexagon} \\ \diagup \\ x_1 \\ \end{array} \\ &= \text{Li}_{k;1,1}^{\mathcal{L}} ([x_1, x_2, x_3, x_6], [x_3, x_4, x_5, x_6]) - \text{Li}_{k;1,1}^{\mathcal{L}} ([x_1, x_2, x_5, x_6], [x_3, x_4, x_5, x_2]) \\ &\quad + \text{Li}_{k;1,1}^{\mathcal{L}} ([x_1, x_4, x_5, x_6], [x_1, x_2, x_3, x_4]) \end{aligned}$$


General functional equation

$$\mathbf{Q}_{2k}: \quad \sum_{i=1}^{2k+3} (-1)^i \text{QLi}_{2k}(x_1, \dots, \hat{x}_i, \dots, x_{2k+3}) \equiv 0 \pmod{\text{depth} < k}$$

Ingredients (II) – Stable genus 0 curves

Part of boundary strata of Deligne-Mumford compactification $\overline{\mathcal{M}}_{0,n}$
 [[From Hannah Larson's talk, Monday]]

- Components are isomorphic to \mathbb{P}^1
- Only singular points are simple double points
- Number of marked & singular points per components ≥ 3



Gadget to encode limits of projective configurations

Idea: The points x_1, x_3, x_8 all go to p


But: There is always a projective transformation moving (x_1, x_3, x_8, p) to $(\infty, 0, 1, z)$

So: Points x_1, x_3, x_8 split off as a separate \mathbb{P}^1

Cross-ratios well-defined: $[x_1, x_3, x_8, x_i] = [x_1, x_3, x_8, p]$ and $[x_1, x_3, x_2, x_4] = [x_1, x_3, p, p] = 0$

Ingredients (III) – Build to symmetries step-by-step

Specialise \mathbf{Q}_6 to



and use $\lim_{x_i \rightarrow 0 \text{ or } \infty} \text{Li}_{k_1, \dots, k_d}^{\mathcal{L}}(x_1, \dots, x_d) = 0 \pmod{dp < d}$

Write $f(x, y, z) := \text{Li}_{4,1,1}^{\mathcal{L}}((xyz)^{-1}, y, z)$, as before

Lemma (Full Symmetry 1)

$$f(A, B, C) \equiv -f\left(1 - A, \frac{B}{B - 1}, 1 - C\right) \pmod{\text{depth } 2}$$

Remark: First needed to show $f(1, x, y) \equiv 0 \pmod{\text{depth } 2}$, already non-trivial(!)

- Find a second *different* symmetry, and a four-term relation
- Play the symmetries against each other
- Extract (eventually...) the Zagier-type reduction

Summary

- Depth reductions of MPL's
 - Appearances in Number Theory, Hyperbolic Geometry, High-Energy Physics
- Goncharov's Depth Conjecture
 - Cobracket on (motivic) MPL's
 - Truncation and iteration of the cobracket
 - Cobracket conjecturally detects depth
- Zagier-, & Gangl-type reductions in weight 4, and weight 6
- Ideas from the proof
 - Quadrangular polylogarithm functional equations
 - Degenerations to stable genus 0 curves
 - Discover and exploit many symmetries/short identities